

Transmission between International Markets & Causality

A. Charpentier (Université de Rennes 1)



Discussion of

*Testing for Extreme Volatility Transmission
with Realized Volatility Measures*

By Sessi Tokpavi,

with Christophe Boucher, Gilles de Truchis and Elena Dumitrescu

IMAC3, International MACroeconomics Workshop, December 2016.

<http://freakonometrics.hypotheses.org>

Granger Causality

Causality $2 \rightarrow 1$,

$$\mathbb{P}[X_{t+1}^{(1)} \in \mathcal{A} \mid \mathcal{F}_t^{(1,2)}] \neq \mathbb{P}[X_{t+1}^{(1)} \in \mathcal{A} \mid \mathcal{F}_t^{(1)}]$$

see [Granger \(1969\)](#), or more conveniently

$$f(x_{t+1}^{(1)} \mid x_t^{(1)}) \neq f(x_{t+1}^{(1)} \mid x_t^{(1)}, x_t^{(2)}).$$

Classical interpretation on the first moment

$$\mathbb{E}[X_{t+1}^{(1)} \mid \mathcal{F}_t^{(1,2)}] \neq \mathbb{E}[X_{t+1}^{(1)} \mid \mathcal{F}_t^{(1)}]$$

with a linear model (VAR) it is a nullity test, i.e.

$$X_{t+1}^{(1)} = \phi_{11} X_t^{(1)} + \underbrace{\phi_{12}}_{\substack{? \\ =0}} X_t^{(2)} + \varepsilon_{t+1}$$

See also [Taamouti et al. \(2012\)](#) or [Bouezmarni et al. \(2009\)](#).

Volatility transmission between international markets

Volatility transmission with High Frequency Data

Use of **realized volatility** to estimate the true latent process of volatility

But as mentioned in **Corradi, Distaso & Fernandes (2012)** it is necessary to control for the effect of jumps and microstructure noise in the log-price process when testing for spillover effects in the integrated variance.

Here volatility transmission test focuses only on extreme or large values of the volatility process.

The Model

Underlying **continuous time model**

$$d\mathbf{p}(t) = \boldsymbol{\mu}(t)dt + \boldsymbol{\Sigma}(t)d\mathbf{W}_t + \boldsymbol{\xi}d\mathbf{N}_t$$

with $\mathbf{J}(t) = \boldsymbol{\xi}(t)d\mathbf{N}_t$.

From an econometrician perspective, we have **discrete time observed return**

T = number of days and $M = \Delta^{-1}$ = number of intraday returns,

$$\mathbf{r}_{t+j\Delta} = \mathbf{p}_{t+j\Delta}^* - \mathbf{p}_{t+(j-1)\Delta}^* \text{ where } \mathbf{p}_{t+j\Delta}^* = \mathbf{p}_{t+j\Delta} + \boldsymbol{\epsilon}_{t+j\Delta}$$

Realized variance (see **Barndorff-Nielsen & Shephard (2001)**) is

$$\mathbf{RV}_{t,M} = \sum_{j=1}^M \mathbf{r}_{t-1+j\Delta} \mathbf{r}_{t-1+j\Delta}^\top$$

Inference

Quadratic variation is $QV_{t+1} = \underbrace{\int_t^{t+1} \Sigma(\tau) d\tau}_{IV_{t+1}} + \underbrace{\sum_{\tau=t}^{t+1} J(\tau)J(\tau)^\top d\tau}_{JV_{t+1}}$ A consistent

nonparametric estimator of QV_{t+1} (see [Barndorff-Nielsen & Shephard \(2002\)](#)) is $RV_{t,M}$, since $RV_{t,M} \xrightarrow{\mathbb{P}} Q_t$ as $M \rightarrow \infty$.

A consistent nonparametric estimator of IV_{t+1} (see [Barndorff-Nielsen & Shephard \(2002\)](#)) is $BV_{t,M}$ (bipolar variation), with

$$BV_{t,M} = \frac{\pi}{2} \sum_{j=2}^M |\mathbf{r}_{t-1+j\Delta}| |\mathbf{r}_{t-1+(j-1)\Delta}^\top|$$

IV_t = integrated variance, with cdf $F_t(\cdot)$, and marginal quantile functions $q_{i,t}(\cdot)$.
Extrem periods are defined as $z_{i,t}(\alpha) = \mathbf{1}(IV_{i,t} > q_{i,t}(\alpha))$.

For non causality $2 \rightarrow 1$, test here $H_0 : \mathbb{E}[z_{1,t}(\alpha) | \mathcal{F}_{t-1}^{(1,2)}] = \mathbb{E}[z_{1,t}(\alpha) | \mathcal{F}_{t-1}^{(1)}]$.

A model for $q_t(\alpha)$

Consider a heterogeneous autoregressive quantile model (HARQ), see [Zikes & Barunik \(2014\)](#)

$$q_t(\alpha) = \theta_1 + \theta_2 \mathbf{IV}_{t-1} + \theta_3 \overline{\mathbf{V}}_t^{(5)} + \theta_4 \overline{\mathbf{V}}_t^{(22)} + \varepsilon_t$$

with $\overline{\mathbf{V}}_t^{(J)} = \frac{1}{J} \sum_{j=1}^J \mathbf{IV}_{t-j}$.

Using [Koenker & Bassett Jr \(1978\)](#), estimate θ_i by solving

$$\operatorname{argmin}_{\theta \in \mathbb{R}^4} \left\{ \frac{1}{T} \sum_{t=2}^T (\alpha - \mathbf{1}(u_{i,t} < 0)) \cdot u_{i,t} \right\} \text{ where } u_{i,t} = \widehat{IV}_{i,t} - q_{i,t}(\alpha, \theta)$$

and then set $\widehat{z}_{i,t}(\alpha) = \mathbf{1}(\widehat{IV}_{i,t} > \widehat{q}_{i,t}(\alpha))$.

Cross-covariance at lag-order h

Then compute cross-covariances

$$\hat{\rho}_h^{(1,2)} = \frac{\hat{C}_h^{(1,2)}}{\alpha(1-\alpha)} \quad \text{with} \quad \hat{C}_h^{(1,2)} = \frac{1}{T-h} \sum_{t=1+h}^T [\hat{z}_{1,t}(\alpha) - (1-\alpha)][\hat{z}_{2,t}(\alpha) - (1-\alpha)]$$

and consider [Ljung & Box \(1978\)](#)'s statistics,

$$\hat{Q}_H^{(1,2)} = T(T+2) \sum_{h=1}^H \frac{[\hat{\rho}_h^{(1,2)}]^2}{T-h}$$

Under H_0 , $\hat{Q}_H^{(1,2)} \sim \xi^2(H)$ as $T, M, \rightarrow \infty$.

From asymptotics to finite sample

$M = 23,400$ (1 second sampling frequency and a trading day of 6.5 hours)

Simulations based on log-volatility follows a stationary three-regime smooth-transition heterogeneous autoregressive (HARST) model